

Whole-blood Gene Expression Profiling in Ankylosing Spondylitis Shows Upregulation of Toll-like Receptor 4 and 5

SHERVIN ASSASSI, JOHN D. REVEILLE, FRANK C. ARNETT, MICHAEL H. WEISMAN, MICHAEL M. WARD, SANDEEP K. AGARWAL, PRAVITT GOURH, JITEN BHULA, ROOZBEH SHARIF, KEERAN SAMPAT, MAUREEN D. MAYES, and FILEMON K. TAN

ABSTRACT. Objective. To identify differentially expressed genes in peripheral blood cells (PBC) of patients with ankylosing spondylitis (AS) relative to healthy controls and controls with systemic inflammation.

Methods. We investigated PBC samples of 16 patients with AS and 14 matched controls, in addition to systemic lupus erythematosus (SLE) and systemic sclerosis (SSc) samples utilizing Illumina Human Ref-8 BeadChips. Candidate genes were confirmed using quantitative PCR. Subsequently, these genes were also validated in a separate sample of 27 patients with AS [before and after anti-tumor necrosis factor (anti-TNF) treatment] and 27 matched controls.

Results. We identified 83 differentially expressed transcripts between AS patients and controls. This gene list was filtered through the lists of differentially expressed transcripts in SLE and SSc, which resulted in identification of 52 uniquely dysregulated transcripts in AS. Many of the differentially expressed genes belonged to Toll-like receptor (TLR) and related pathways. *TLR4* and *TLR5* were the only dysregulated TLR subtypes among AS patients. We confirmed the overexpression of *TLR4* and *TLR5* in AS patients in comparison to controls ($p = 0.012$ and $p = 0.006$, respectively) and SLE ($p = 0.002$, $p = 0.008$) using quantitative PCR in the same sample. Similarly, *TLR4* ($p = 0.007$) and *TLR5* ($p = 0.012$) were significantly upregulated among the AS patients before anti-TNF treatment in the confirmatory sample. *TLR4* ($p = 0.002$) and *TLR5* ($p = 0.025$) decreased significantly after anti-TNF treatment.

Conclusion. PBC gene expression profiling in AS shows an upregulation of *TLR4* and *TLR5*. This supports the importance of TLR subtypes in the pathogenesis of AS that are responsible for the immune response to Gram-negative bacteria. (J Rheumatol First Release Oct 15 2010; doi:10.3899/jrheum.100469)

Key Indexing Terms:

ANKYLOSING SPONDYLITIS TOLL-LIKE RECEPTORS IMMUNE SYSTEM
AUTOIMMUNITY BACTERIA GENE EXPRESSION PROFILING

From the Division of Rheumatology and Immunogenetics, University of Texas Health Science Center at Houston, Houston, Texas; Division of Rheumatology, Cedars-Sinai Medical Center, Los Angeles, California; and NIAMS-National Institutes of Health, Bethesda, Maryland, USA.

Supported by American College of Rheumatology Clinical Investigator Fellowship Award and NIH-KL2RR024149-04 (Dr. Assassi); NIH/NIAMS P01-AR-052915-01 (Dr. Reveille, Dr. Weisman); NIH/CORT P50AR054144 (Dr. Arnett, Dr. Mayes); the Intramural Research Program, NIAMS/NIH (Dr. Ward); and NIH/NCRR Clinical and Translational Sciences Award (CTSA), UL1-RR024148

S. Assassi, MD, MS; J.D. Reveille, MD; F.C. Arnett, MD, Division of Rheumatology and Immunogenetics, University of Texas Health Science Center at Houston; M.H. Weisman, MD, Division of Rheumatology, Cedars-Sinai Medical Center; M.M. Ward, MD, MPH, NIAMS-National Institutes of Health; S.K. Agarwal, MD, PhD; P. Gourh, MD; J. Bhula, BS; R. Sharif, MD; K. Sampat, MD; M.D. Mayes, MD, MPH; F.K. Tan, MD, PhD, Division of Rheumatology and Immunogenetics, University of Texas Health Science Center at Houston.

Address correspondence to Dr. S. Assassi, Department of Medicine, Division of Rheumatology, University of Texas-Houston, 6431 Fannin, MSB 5.270, Houston, TX 77030. E-mail: shervin.assassi@uth.tmc.edu
Accepted for publication August 21, 2010.

Ankylosing spondylitis (AS) is a chronic inflammatory arthritis with a predilection for the spine and sacroiliac joints that can lead to new bone formation and ultimately ankylosis. AS is the prototype of spondyloarthropathies (SpA), a related family of disorders with common clinical features and with a strong association with HLA-B27. However, HLA-B27 accounts only for ~45% of the genetic risk in AS. Genome-wide association studies have identified several other non-HLA susceptibility genes such as IL23R and ERAP1 in AS^{1,2}. Other diseases belonging to the spectrum of SpA are reactive arthritis, psoriatic arthritis, and arthritis in patients with inflammatory bowel disease³.

Functional studies also have been undertaken to identify candidate genes and pathways that play a role in the pathogenesis of SpA. Microarray data from synovium suggest a proinflammatory profile. Gu, *et al* demonstrated increased RNA expression of monocyte chemoattractant protein 1

(MCP-1), interleukin 8 (IL-8), IL-1 β , endothelial-monocyte activating polypeptide II, interferon- γ , tumor necrosis factor- α (TNF- α), and BiP in SpA synovial fluid mononuclear cells⁴. Rihl, *et al* found elevated levels of IL-7 transcript and protein in sacroiliac joint cells⁵. On the other hand, transcriptional profiling of isolated peripheral blood mononuclear cells (PBMC) by several groups showed that transcripts involved in the inflammatory response are differentially expressed in SpA patients, but reports on the nature of these changes seem to vary. The earliest PBMC study indicated increased expression of proinflammatory proteins such as CXCR4/SDF-1⁶. However, recent reports suggest decreased immune responsiveness of the PBMC. Smith, *et al* found a “reverse interferon (IFN) signature” characterized by decreased expression of IFN- γ and IFN- γ -inducible genes in AS macrophages⁷, and Duan, *et al* found that AS PBMC display an immunosuppressive phenotype as shown by underexpression of NR4A2, TNFAIP3, and CD69⁸.

We investigated the whole-blood gene transcript profile of AS patients in comparison to controls in order to elucidate the gene expression patterns involved in this disease. Unlike previous investigators, we employed a commercially available method to stabilize RNA in the blood immediately upon phlebotomy to minimize artifacts that occur with handling and purification of living PBMC⁹. We compared their transcript profiles to healthy controls; we also compared transcript profiles to those of patients with systemic sclerosis (SSc) and systemic lupus erythematosus (SLE) in order to identify transcripts that were specific to AS and were not related only to the presence of systemic inflammation. We identified 51 genes that were differentially expressed only in AS patients. Many of the differentially expressed genes belonged to Toll-like receptor (TLR) and related pathways. We observed an overexpression of *TLR4* and *TLR5* that was confirmed by quantitative polymerase chain reaction (PCR) in 2 different cohorts of AS patients. We further demonstrated that *TLR4* and *TLR5* transcripts in whole blood decreased significantly after TNF- α inhibitor (anti-TNF) therapy. Our findings provide support for the importance of a pathogen-associated molecular pattern in the pathogenesis of AS.

MATERIALS AND METHODS

The patients with AS were recruited from the Prospective Study of Outcomes in Ankylosing Spondylitis (PSOAS) study. The PSOAS is a longitudinal study of AS patients from 3 sites in the USA: the University of Texas Health Science Center at Houston, Houston, Texas (UTHSC-H); Cedars-Sinai Medical Center, Los Angeles, California; and the National Institutes of Health, Bethesda, Maryland. All AS patients met the modified New York criteria for the definitive diagnosis of AS¹⁰. From patients enrolled in PSOAS, we investigated 2 separate groups of patients with AS. In the initial group, no patient was receiving anti-TNF or other immunosuppressive agents. All AS patients in the initial sample set were recruited from the UTHSC-H site. In the second confirmatory cohort, 2 samples were investigated from each patient: the first before initiation of an anti-TNF treatment; the second was obtained after the patient was treated with an

anti-TNF agent for ≥ 6 months. AS patients in the confirmatory cohort were recruited from all 3 participating sites. All AS patients enrolled in the discovery group had active disease, defined as Bath AS Disease Activity Index (BASDAI) score ≥ 3.5 . Similarly, all AS patients in the confirmatory cohort had a BASDAI ≥ 3.5 before the initiation of anti-TNF therapy. In addition to demographic information and BASDAI, we measured C-reactive protein (CRP) at the time of blood draw in AS patients.

The healthy controls had no history of autoimmune diseases or spondyloarthritis-related manifestations and were matched for sex, age, and ethnicity to AS patients. Patients with SLE and SSc were also investigated as disease controls in order to identify gene expression patterns that are specific to AS and are not related only to presence of systemic inflammation. Patients with SSc or SLE were recruited from the continuing longitudinal studies or clinical practice of the investigators at UTHSC-H. All SSc patients met the 1980 American College of Rheumatology (ACR) preliminary criteria for the classification of SSc¹¹. Similarly, all SLE patients fulfilled the ACR classification criteria for SLE¹² and had signs of active disease in at least 2 categories of the Systemic Lupus Activity Measure-Revised¹³. Patients with SSc or SLE receiving immunosuppressive agents other than low-dose steroids (prednisone ≤ 5 mg) and hydroxychloroquine were excluded from the study. The comparison group for the patients with SSc or SLE were healthy controls matched for sex, age, and ethnicity to patients with SSc.

All study subjects provided written informed consent and the study was approved by the institutional review boards of all participating centers.

Sample processing and microarray experiments. Whole-blood samples for gene expression studies were drawn directly into PAXgeneTM tubes (PreAnalytix, Hombrechtikon, Switzerland). All blood samples were processed in the laboratories of the Division of Rheumatology and Clinical Immunogenetics, UTHSC-H. Total RNA was isolated and purified according to the manufacturer’s protocol using PAXgene RNA Kit. The RNA quality and yield was assessed by NanoDrop ND-1000 Spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA) and Agilent 2100 Bioanalyzer (Agilent Technologies, Palo Alto, CA, USA). A globin reduction was not done because this procedure did not increase percentage of present calls in a preliminary experiment with 9 healthy control samples. This finding might be explained by the longer transcript probes printed on the Illumina arrays (50 mer probes) in comparison to Affymetrix arrays.

Two hundred nanograms of total RNA were amplified and purified utilizing Illumina TotalPrep RNA Amplification Kit (Applied Biosystems/Ambion, Austin, TX, USA) according to the manufacturer’s instructions. We hybridized the amplified cRNA on Illumina Human Ref-8v2 arrays and extracted the data utilizing the Illumina Beadstudio software (Illumina, San Diego, CA, USA).

Microarray data analysis. The raw data were exported into BRB-ArrayTools v. 3.7 (R. Simon and A. Pen Lam, National Cancer Institute, Bethesda, MD, USA).

Probes whose signal detection p values indicated no significant difference from those of the negative controls ($p < 0.01$) were removed from the analysis. In addition, we excluded genes whose expression values were missing or were filtered out in more than 50% of experiments. Expression data were normalized using the median over the entire array. We used less stringent criteria for detection of differentially expressed genes in order to increase detection of any transcripts with altered gene expression. A gene was defined as differentially expressed for all comparisons when the significance level for comparison was $p < 0.01$, utilizing a random variance t test¹⁴. The set of differentially expressed genes was modeled in Ingenuity Pathway Analysis (Ingenuity Systems, Redwood City, CA, USA) to detect pathways or biological processes involving these genes.

Real-time quantitative PCR. Quantitative PCR (qPCR) assays for *TLR4* and *TLR5* were designed to confirm the microarray results. The assay details including the primer sequence, lowest limit of detection, and PCR efficiency are shown in Table 1. Each sample was assayed in triplicate plus a control without reverse transcriptase to access DNA contamination. Samples

Table 1. Details of quantitative PCR.

Gene	Accession No.	Primer and Probe	Amplicon Length	PCR Efficiency	Lower Limit of Detection (molecules)
Toll-like receptor 4	NM_138554	500 (+) GAG CCT TTT CTG GAC TAT CAA G* 582 (-) TCC AAT GGG GAA GTT CTC TAG* 554 (-) FAM-AGA TTT GTC TCC ACA GCC ACC AGC-BHQ1†	81 bases	95%	180
Toll-like receptor 5	NM_003268	2209 (+) GCC ATC TGA CTG CAT TAA GG* 2284 (-) GCA GGT AAA TCA TTG TGA GAA AG* 2237 (+) FAM-CCT CAA CTC CAA CAG GCT GAC AGT-BHQ1†	84 bases	95%	170

* Primer sequence. † Probe sequence.

were reverse transcribed into cDNA using superscript II reverse transcriptase (Invitrogen, Carlsbad, CA, USA) for 30 min at 50°C in 384-well plates. PCR master mix containing JumpStart Taq Polymerase (Sigma, St. Louis, MO, USA) was added to the samples. Each assembled plate was run in a 7900 real-time instrument using the following PCR conditions: 95°C for 1 min, followed by 40 cycles of 95°C for 12 s and 60°C for 30 s. Results were analyzed utilizing SDS 2.3 (7900) software (Applied Biosystems, Foster City, CA, USA) with FAM reporter and ROX as the reference dye. The final data were normalized to 18s rRNA levels. The final data were presented as the molecules of the transcript divided by the molecules of 18sRNA transcript × 100.

Statistical analysis. Continuous variables were analyzed by t test if the dependent variable had a normal distribution. The Mann-Whitney nonparametric test was used if the outcome variable did not have a normal distribution. We compared the gene expression values, CRP, and BASDAI scores between pre- and post-anti-TNF treatment samples utilizing a paired T test if model assumptions were met, otherwise Wilcoxon signed-rank test was applied. Linear regression was used to investigate the relationship among 2 continuous variables. Two-sided p values < 0.05 were considered significant. Analyses were performed utilizing the NCSS 2007 statistical program (NCSS, Kaysville, UT, USA).

RESULTS

Characteristics of study groups. A total of 16 patients with AS and 14 healthy controls were examined in the first group. There were no significant differences in age, sex, and ethnicity between patients and controls. Table 2 shows the demographic characteristics, presence of spondyloarthritis (SpA) related manifestations, and other clinical features in

participants. No control subjects had a SpA-related manifestation. The majority of AS patients (75%) were male. Among AS patients, the mean BASDAI score was 5.31 (± 2.01 SD) and the median CRP was 0.59 mg/dl. All AS patients in this group were HLA-B27-positive. We also investigated the gene expression profile of 74 patients with SSc (female 79.7%, mean age 49.16 yrs), 21 matched controls (female 80.95%, mean age 53.53 yrs), and 17 patients with SLE (female 94.12%, mean age 38.5 yrs).

Gene expression microarrays show upregulation of nuclear factor-κB and TLR pathways in AS. A total of 8230 transcripts passed our filtering criteria across all the peripheral blood samples. Clustering analyses according to date of hybridization or chip number indicated no technical artifact.

A comparison of 16 AS patients with their matched controls revealed 83 differentially expressed transcripts. An unsupervised hierarchical clustering of these genes in AS patients and their controls is shown in Figure 1. A list of these genes is provided in Appendix 1. In a similar analysis, comparison of SLE and SSc samples to their matched controls resulted in 936 and 530 differentially expressed transcripts, respectively. The most prominent gene expression pattern among SLE and SSc patients consisted of upregulation of genes belonging IFN-related pathways¹⁵, whereas we did not observe an overrepresentation of up- or down-

Table 2. Demographic features of patients with AS and their matched controls in both samples.

Feature	Discovery Group		Confirmation Group	
	AS, n = 16	Control, n = 14	AS, n = 27	Control, n = 27
Age, mean (SD) yrs	36.01 (11.94)	32.11 (7.34)	36.08 (11.4)	37.21 (10.68)
No. male (%)	12 (75)	10 (71.43)	14 (51.85)	14 (51.85)
No. Caucasian (%)	13 (81.25)	11 (78.57)	17 (62.96)	17 (62.96)
Disease duration	16.09 (12.87)	NA	11.55 (5.86)	NA
BASDAI*	5.31 (2.01)	NA	5.93 (1.35)	NA
C-reactive protein	1.4 (2.23)	NA	1.56 (2.17)	NA
Uveitis	3 (18.75)	0	14 (51.85)	0
Psoriasis	1 (6.25)	0	3 (11.11)	0
Peripheral inflammatory arthritis	10 (62.5)	0	7 (25.93)	0
Crohn's disease	0	0	3 (11.11)	0

BASDAI: Bath Ankylosing Spondylitis Disease Activity Index; NA: not applicable.




Figure 1. Unsupervised hierarchical clustering of 83 genes identified as differentially expressed between patients with AS and controls.

regulated genes belonging to IFN pathways in comparisons of AS patients to controls in the Ingenuity Pathway Analysis.

Then we filtered the differentially expressed genes in AS through the lists of differentially expressed transcripts in SLE and SSc, which resulted in 52 transcripts corresponding to 51 genes that were uniquely differentially expressed among AS patients (Appendix 2). Ingenuity Pathway Analysis of these 52 transcripts demonstrated overrepresentation of transcripts belonging to pathways involved in TLR signaling. *TLR4* ($p = 0.008$) and *TLR5* ($p = 0.006$) were overexpressed in AS patients but not in patients with SLE or SSc. No other subtype of *TLR* was differentially expressed in AS patients.

The significantly dysregulated pathways were nuclear factor- κ B signaling, dendritic cell maturation, TLR, TREM1 signaling, and BRCA1 in DNA damage (Table 3). It is notable that TLR play a major role in the activation of the first 4 pathways. Specifically, *TLR4* and *TLR5* were the only dysregulated genes that were present in all these 4 pathways.

We next assessed *TLR4* and *TLR5* levels in patients with AS, their matched controls, and patients with SLE using qPCR. In agreement with the microarray results, AS patients had higher *TLR4* ($p = 0.012$) and *TLR5* ($p = 0.006$) levels than controls. Similarly, in comparison to SLE, AS patients showed overexpression of *TLR4* ($p = 0.002$) and *TLR5* ($p = 0.008$), whereas *TLR4* ($p = 0.203$) and *TLR5* ($p = 0.383$) levels did not differ significantly between SLE patients and controls. Figure 2 shows the *TLR4* and *TLR5* levels among these 3 study groups. Further, *TLR4* and *TLR5* levels correlated highly with each other ($p < 0.001$, $r^2 = 0.62$).

Overexpression of TLR4 and TLR5 was confirmed in a separate group of AS patients. We next examined the *TLR4* and *TLR5* levels in a second larger sample of 27 patients with AS and 27 matched healthy controls. A total of 22 patients with AS (81.5%) were HLA-B27-positive; characteristics of patients and controls in the confirmation group are shown in Table 2. One sample before and another one after anti-TNF treatment were investigated on each AS patient. The CRP and BASDAI score of AS patients before anti-TNF treatment were 0.791 mg/dl (median) and 5.93 ± 1.35 (mean \pm SD), respectively. As expected, both CRP ($p = 0.005$) and BASDAI score ($p < 0.001$) decreased significantly upon anti-TNF treatment to 0.095 mg/dl (median) and 3.49 ± 2.33 (mean \pm SD), respectively.

Table 3. Dysregulated pathways in comparison of patients with AS and unaffected controls according to Ingenuity Pathway Analysis.

Canonical Pathways	P
Nuclear factor- κ B signaling	0.0006
Dendritic cell maturation	0.008
Toll-like receptor signaling	0.008
TREM1 signaling	0.01
BRCA1 in DNA damage response	0.009

Compared to their matched controls, we again observed higher *TLR4* ($p = 0.007$) and *TLR5* ($p = 0.012$) levels in AS patients before initiation of the anti-TNF treatment. However, *TLR4* and *TLR5* levels in AS samples after anti-TNF treatment did not differ significantly from controls ($p = 0.126$, $p = 0.173$, respectively). Figure 3 shows *TLR4* and *TLR5* levels in the AS samples before and after initiation of anti-TNF treatment and in controls. Similarly to the initial sample, *TLR4* and *TLR5* levels also correlated significantly with each other ($p < 0.001$, $r^2 = 0.44$).

TLR4 and TLR5 levels decreased significantly after TNF blockade. *TLR4* levels decreased significantly after initiation of anti-TNF treatment from 0.072 to 0.046 ($p = 0.002$) in patients with AS. Similarly, the *TLR5* levels also declined significantly on treatment with anti-TNF from 0.005 to 0.0038 ($p = 0.025$; Figure 3). Further, the percentage changes in *TLR4* and *TLR5* after TNF blockade correlated highly with each other ($p < 0.001$, $r^2 = 0.75$).

TLR4 and TLR5 levels in relationship with clinical features. BASDAI scores did not correlate with *TLR4* and *TLR5* levels in either study cohort (data not shown). Although CRP levels correlated with *TLR4* and *TLR5* levels in the initial sample ($p = 0.015$, $r^2 = 0.4$ and $p = 0.001$, $r^2 = 0.6$, respectively), we could not confirm this finding among pre-TNF samples in the second study group ($p = 0.833$ for *TLR4*, $p = 0.753$ for *TLR5*). Among the patients with AS, history of uveitis, psoriasis, peripheral inflammatory arthritis, or Crohn's disease did not correlate with *TLR4* and *TLR5* levels in the 2 study cohorts (data not shown). Further, there was no significant difference in *TLR4* and *TLR5* levels between the HLA-B27-positive and HLA-B27-negative AS patients ($p = 0.454$, $p = 0.319$, respectively).

DISCUSSION

We observed that PBC global gene expression profiling of patients with AS showed a dysregulation of TLR-related pathways. We specifically identified an overexpression of *TLR4* and *TLR5*, which also was confirmed by qPCR in the initial and a separate confirmatory sample. Differential regulation of these genes appeared to be unique to AS and was not observed in SLE or SSc. Our study is the first report of increased expression of *TLR5* in AS.

An abnormal host response against pathogens has been implicated in the pathogenesis of AS and other SpA subtypes. Further, 60% of patients with SpA without evidence of clinical Crohn's disease have endoscopic or histological signs of gut inflammation¹⁶. Moreover, studies with B27-transgenic rats provide support for the role of commensal gut flora in the pathogenesis of HLA-B27-associated gut and joint manifestations. The B27-transgenic rats do not develop inflammatory intestinal or peripheral joint disease in a germ-free environment¹⁷.

TLR are primarily involved in innate immune responses to microbial pathogens by recognition of conserved pathogen-




Figure 2. Gene expression levels of *TLR4* and *TLR5* in AS, SLE, and controls by qPCR in the first study sample. Expression of both genes is higher in AS patients in comparison to controls and SLE patients, while there was no significant difference between SLE patients and controls.




Figure 3. Gene expression levels of *TLR4* and *TLR5* in patients with AS before and during anti-TNF treatment, as well as controls, in the second study sample. Gene expression levels of both genes decreased significantly after anti-TNF treatment among AS patients. Again, before anti-TNF treatment, samples showed higher *TLR4* and *TLR5* levels than controls. After anti-TNF treatment, there was no significant difference between AS patients and controls.

associated molecular patterns^{18,19}. More than 10 TLR subtypes have been identified. In our study, *TLR4* and *TLR5* were the only TLR subtypes that were overexpressed among patients with AS. Lipopolysaccharide (LPS) in the outer membrane of Gram-negative bacteria is the main ligand of *TLR4*¹⁹. In animal models, *TLR4* plays a critical role in the early cytokine response of phagocytes upon infection with reactive arthritis-associated Gram-negative bacteria such as *Yersinia*, *Salmonella*, and *Chlamydia*^{20,21,22}. An upregulation of *TLR4* among patients with AS has been previously reported. De Rycke, *et al* reported increased expression of *TLR4* but not *TLR2* on PBMC of patients with SpA in comparison to controls²³. Yang, *et al* showed that the expression of *TLR4* on lymphocytes, monocytes, and neutrophils was all significantly increased among patients with AS²⁴.

The main ligand for *TLR5* is flagellin, a primary component of bacterial flagella that extend from the outer mem-

brane of Gram-negative bacteria. Flagella are known to be major antigens of Gram-negative bacteria like *Salmonella*, *Escherichia coli*, and *Yersinia*, where its antigenicity serves as the basis for H serotyping^{25,26,27}. There are no published reports on the role of *TLR5* in AS or other SpA subtypes. However, both *TLR5* and flagellin have been implicated in the pathogenesis of Crohn's disease (reviewed by Gewirtz²⁸). A study investigating the proteins of commensal microflora that were reactive with antisera from a colitic mouse model identified flagellins as the dominant antigen. Further, serum IgG to these flagellins was elevated in patients with Crohn's disease but not in patients with ulcerative colitis or controls²⁹. In another study, flagellin exposure to injured mouse colon *in vivo*, but not intact colon, significantly worsened colonic inflammation, whereas *TLR2*-specific agonists did not have a similar effect³⁰. A *TLR5* stop-polymorphism was negatively associated with

Crohn's disease in Ashkenazi Jewish patients, raising the possibility that a downregulation in *TLR5* provides protection against development of Crohn's disease³¹.

AS and Crohn's disease share common clinical features and genetic background. Up to one-third of patients with Crohn's disease have sacroiliac joint involvement similar to AS on computer tomography³², whereas roughly 60% of patients with SpA have subclinical colitis by biopsy¹⁶. These similarities, along with our data, suggest that the role of *TLR5* in the pathogenesis of AS should be further explored in mechanistic studies. We did not observe any significant difference in *TLR5* levels between the AS patients with and those without Crohn's disease in our study ($p = 0.769$). This finding suggests that the overexpression of *TLR5* in AS is independent of the presence of clinically apparent Crohn's disease. However, our study might have been underpowered to detect a relationship between *TLR5* expression and presence of Crohn's disease in patients with AS.

Activation of the innate immune system such as polymorphonuclear cells and macrophages plays an important role in AS-related inflammation. Patients with SpA have significantly higher neutrophil counts and lower lymphoid aggregates than patients with rheumatoid arthritis (RA) on synovial histology³³. Further, macrophages expressing the hemoglobin scavenger receptor CD163 are increased in the synovial lining of patients with SpA compared to RA patients. CD163+ macrophages also are increased in the colonic lamina propria in SpA patients compared to controls, supporting the hypothesis of a recirculation of similar clones in the intestinal mucosa and synovium³⁴. Of interest, both *TLR4* and *TLR5* can induce an acute shedding of CD163 from human monocytes. As well, these 2 TLR subtypes have a synergistic effect on upregulation of CD163, whereas exogenous recombinant IFN- γ leads to downregulation of CD163³⁵. Global gene expression studies of macrophages derived from AS patients reveal a "reverse" IFN signature, with IFN- γ upregulated genes being downregulated⁷. The observed upregulation of *TLR4* and *TLR5* in our study, along with the reported "reverse" IFN- γ in AS, are both potential mechanisms that could lead to overexpression CD163+ macrophages in this disease. However, this hypothesis needs to be verified by further mechanistic studies. We did not observe a "reverse" IFN signature in whole-blood samples as described by Smith, *et al* in macrophages of patients with AS⁷.

We observed downregulation of *TLR4* and *TLR5* after initiation of anti-TNF treatment in patients with AS. Both *TLR4* and *TLR5* can induce secretion of TNF- α and other proinflammatory cytokines^{30,36}. The expression of *TLR4* on PBMC decreases gradually after treatment of SpA patients with anti-TNF agents *in vivo*. These PBMC have a functional impairment in their capacity to produce TNF- α after stimulation with LPS *in vitro*²³. Further, *TLR4* mRNA correlated closely with serum TNF- α levels among patients

with AS²⁴. These findings suggest that TNF blockade attenuates a self-perpetuating activation of the innate immune system via the TLR pathway.

The *TLR4* and *TLR5* levels and their percentage changes after TNF blockade correlated closely with each other, which raises the possibility that they are both triggered by the same mechanism, whereas we did not observe such a strong correlation of these 2 transcripts with general markers of inflammation such as CRP.

The data on correlation of *TLR4* and *TLR5* levels with CRP in our study were not consistent. While we observed a significant correlation in the first sample, we could not confirm this finding in the second cohort. These results did not change even after we excluded the HLA-B27-negative patients from the analysis (data not shown). Similarly, De Rycke, *et al*²³ did not observe a correlation of *TLR4* expression with CRP in SpA, whereas Yang, *et al*²⁴ reported correlation of *TLR4* mRNA levels with CRP in HLA-B27-positive patients with AS. Studies with larger sample sizes are needed to resolve this issue.

Potential limitations of our study are that we used less stringent criteria for identification of differentially expressed genes in microarray data analysis in order to increase our ability to detect dysregulated genes and pathways. However, we verified the overexpression of *TLR4* and *TLR5* with qPCR in the same sample in addition to a separate confirmatory cohort. We compared the global gene expression of AS patients to transcriptomes of patients with SLE and SSc. These comparative studies should be extended to patients with other rheumatological diseases such as RA in future investigations. Further, the observed fold-changes in *TLR4* and *TLR5* between AS patients and controls were relatively small (< 2-fold). However, whole-blood samples consist of heterogeneous populations of white blood cell subtypes. It is possible the observed differential expression levels in whole blood are secondary to much higher-fold changes in a particular subset of white blood cells. We focused on dysregulation of TLR transcripts. However, the other observed differentially expressed transcripts and pathways in this study also could play an important role in the pathogenesis of AS. For example, the observed dysregulations in the dendritic cell maturation pathway may provide further support for the importance of defective functional capacity of dendritic cells in SpA³⁷.

In summary, our global gene expression analysis revealed the TLR-related pathways as the most prominently dysregulated biological process in AS. We identified *TLR4* and *TLR5* as the only dysregulated subtypes of TLR in AS. We confirmed the overexpression of these 2 genes among patients with AS in the same sample and in a confirmatory cohort. We showed that the expression of both receptors decreased after initiation of anti-TNF treatment. Our findings provide further support for the importance of TLR subtypes responsive to Gram-negative bacteria in the pathogen-

esis of AS. Mechanistic studies are needed to elucidate the role of these TLR subtypes in the development of AS.

ACKNOWLEDGMENT

The authors thank Laura Diekman, Stephanie Brown, Lori Guthrie, and Vera Wirawan for help with data collection and management. We are also grateful to Dr. David Loose, Dr. Gregory Shipley, Nancy Shipley, and Jun Ying for assistance in design and performance of the laboratory studies.

REFERENCES

1. Burton PR, Clayton DG, Cardon LR, Craddock N, Deloukas P, Duncanson A, et al. Association scan of 14,500 nonsynonymous SNPs in four diseases identifies autoimmunity variants. *Nat Genet* 2007;39:1329-37.
2. Brown MA. Genetics and the pathogenesis of ankylosing spondylitis. *Curr Opin Rheumatol* 2009;21:318-23.
3. Braun J, Sieper J. Ankylosing spondylitis. *Lancet* 2007;369:1379-90.
4. Gu J, Rihl M, Marker-Hermann E, Baeten D, Kuipers JG, Song YW, et al. Clues to pathogenesis of spondyloarthropathy derived from synovial fluid mononuclear cell gene expression profiles. *J Rheumatol* 2002;29:2159-64.
5. Rihl M, Kellner H, Kellner W, Barthel C, Yu DT, Tak PP, et al. Identification of interleukin-7 as a candidate disease mediator in spondylarthritis. *Arthritis Rheum* 2008;58:3430-5.
6. Gu J, Marker-Hermann E, Baeten D, Tsai WC, Gladman D, Xiong M, et al. A 588-gene microarray analysis of the peripheral blood mononuclear cells of spondyloarthropathy patients. *Rheumatology* 2002;41:759-66.
7. Smith JA, Barnes MD, Hong D, DeLay ML, Inman RD, Colbert RA. Gene expression analysis of macrophages derived from ankylosing spondylitis patients reveals interferon-gamma dysregulation. *Arthritis Rheum* 2008;58:1640-9.
8. Duan R, Leo P, Bradbury L, Brown MA, Thomas GP. Gene expression profiling reveals a downregulation in immune-associated genes in patients with AS. *Ann Rheum Dis* 2010;69:1724-9.
9. Baechler EC, Batliwalla FM, Karypis G, Gaffney PM, Moser K, Ortmann WA, et al. Expression levels for many genes in human peripheral blood cells are highly sensitive to ex vivo incubation. *Genes Immun* 2004;5:347-53.
10. Goie The HS, Steven MM, van der Linden SM, Cats A. Evaluation of diagnostic criteria for ankylosing spondylitis: a comparison of the Rome, New York and modified New York criteria in patients with a positive clinical history screening test for ankylosing spondylitis. *Br J Rheumatol* 1985;24:242-9.
11. Subcommittee for Scleroderma Criteria of the American Rheumatism Association Diagnostic and Therapeutic Criteria Committee. Preliminary criteria for the classification of systemic sclerosis (scleroderma). *Arthritis Rheum* 1980;23:581-90.
12. Tan EM, Cohen AS, Fries JF, Masi AT, McShane DJ, Rothfield NF, et al. The 1982 revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum* 1982;25:1271-7.
13. Uribe AG, Vila LM, McGwin G Jr, Sanchez ML, Reveille JD, Alarcon GS. The Systemic Lupus Activity Measure-revised, the Mexican Systemic Lupus Erythematosus Disease Activity Index (SLEDAI), and a modified SLEDAI-2K are adequate instruments to measure disease activity in systemic lupus erythematosus. *J Rheumatol* 2004;31:1934-40.
14. Wright GW, Simon RM. A random variance model for detection of differential gene expression in small microarray experiments. *Bioinformatics* 2003;19:2448-55.
15. Assassi S, Mayes MD, Arnett FC, Gourh P, Agarwal SK, McNearney TA, et al. Systemic sclerosis and lupus: points in an interferon-mediated continuum. *Arthritis Rheum* 2010;62:589-98.
16. Mielants H, Veys EM, Cuvelier C, De Vos M, Goemaere S, De Clercq L, et al. The evolution of spondyloarthropathies in relation to gut histology. II. Histological aspects. *J Rheumatol* 1995;22:2273-8.
17. Taurog JD, Richardson JA, Croft JT, Simmons WA, Zhou M, Fernandez-Sueiro JL, et al. The germfree state prevents development of gut and joint inflammatory disease in HLA-B27 transgenic rats. *J Exp Med* 1994;180:2359-64.
18. Miyake K. Innate immune sensing of pathogens and danger signals by cell surface Toll-like receptors. *Semin Immunol* 2007;19:3-10.
19. Takeda K, Kaisho T, Akira S. Toll-like receptors. *Annu Rev Immunol* 2003;21:335-76.
20. Bulut Y, Faure E, Thomas L, Karahashi H, Michelsen KS, Equils O, et al. Chlamydial heat shock protein 60 activates macrophages and endothelial cells through Toll-like receptor 4 and MD2 in a MyD88-dependent pathway. *J Immunol* 2002;168:1435-40.
21. Sing A, Tvardovskaia N, Rost D, Kirschning CJ, Wagner H, Heesemann J. Contribution of toll-like receptors 2 and 4 in an oral *Yersinia enterocolitica* mouse infection model. *Int J Med Microbiol* 2003;293:341-8.
22. Li Q, Cherayil BJ. Role of Toll-like receptor 4 in macrophage activation and tolerance during *Salmonella enterica* serovar Typhimurium infection. *Infect Immun* 2003;71:4873-82.
23. De Rycke L, Vandooren B, Kruithof E, De Keyser F, Veys EM, Baeten D. Tumor necrosis factor alpha blockade treatment down-modulates the increased systemic and local expression of Toll-like receptor 2 and Toll-like receptor 4 in spondylarthropathy. *Arthritis Rheum* 2005;52:2146-58.
24. Yang ZX, Liang Y, Zhu Y, Li C, Zhang LZ, Zeng XM, et al. Increased expression of Toll-like receptor 4 in peripheral blood leucocytes and serum levels of some cytokines in patients with ankylosing spondylitis. *Clin Exp Immunol* 2007;149:48-55.
25. Johnson JR, Orskov I, Orskov F, Goulet P, Picard B, Moseley SL, et al. O, K, and H antigens predict virulence factors, carboxylesterase B pattern, antimicrobial resistance, and host compromise among *Escherichia coli* strains causing urosepsis. *J Infect Dis* 1994;169:119-26.
26. Aleksic S, Bockemuhl J, Lange F. Studies on the serology of flagellar antigens of *Yersinia enterocolitica* and related *Yersinia* species. *Zentralbl Bakteriol Mikrobiol Hyg A* 1986;261:299-310.
27. Ibrahim GF, Fleet GH, Lyons MJ, Walker RA. Production of potent *Salmonella* H antisera by immunization with polymeric flagellins. *J Clin Microbiol* 1985;22:347-51.
28. Gewirtz AT. TLRs in the gut. III. Immune responses to flagellin in Crohn's disease: good, bad, or irrelevant? *Am J Physiol Gastrointest Liver Physiol* 2007;292:G706-G710.
29. Lodes MJ, Cong Y, Elson CO, Mohamath R, Landers CJ, Targan SR, et al. Bacterial flagellin is a dominant antigen in Crohn disease. *J Clin Invest* 2004;113:1296-306.
30. Rhee SH, Im E, Riegler M, Kokkotou E, O'Brien M, Pothoulakis C. Pathophysiological role of Toll-like receptor 5 engagement by bacterial flagellin in colonic inflammation. *Proc Natl Acad Sci USA* 2005;102:13610-5.
31. Gewirtz AT, Vijay-Kumar M, Brant SR, Duerr RH, Nicolae DL, Cho JH. Dominant-negative TLR5 polymorphism reduces adaptive immune response to flagellin and negatively associates with Crohn's disease. *Am J Physiol Gastrointest Liver Physiol* 2006;290:G1157-G1163.
32. Scott WW Jr, Fishman EK, Kuhlman JE, Caskey CI, O'Brien JJ, Walia GS, et al. Computed tomography evaluation of the sacroiliac joints in Crohn disease. Radiologic/clinical correlation. *Skeletal Radiol* 1990;19:207-10.
33. Kruithof E, Baeten D, De Rycke L, Vandooren B, Foell D, Roth J, et al. Synovial histopathology of psoriatic arthritis, both

Appendix 1. Differentially expressed genes in AS compared to controls.

Parametric p value	Geom mean of intensities in class 1 (AS)	Geom mean of intensities in class 2 (control)	Fold-change	GB acc	Gene symbol	Defined Gene list
0.000478	137.0849484	191.4323799	0.7161012	NM_002340	LSS	Biosynthesis of steroids
0.0006957	9364.319315	4899.472307	1.9112914	NM_000570	FCGR3B	Natural killer cell mediated cytotoxicity, immunology
0.0009312	70.135071	39.2769027	1.7856569	NM_198330	INPP5F	
0.0010988	71.4184127	45.7446495	1.5612408	NM_001255	CDC20	Cell cycle, Ubiquitin mediated proteolysis
0.0011141	52.8656567	70.9770793	0.7448272	NM_001013840	C10orf61	
0.0011586	392.3825626	275.9854116	1.4217511	NM_182755	ZNF438	
0.0017355	61.4036443	44.5498863	1.378312	NM_001125	ADPRH	
0.0018922	87.5243114	47.5085126	1.8422869	NM_170606	MLL3	Lysine degradation
0.002143	53.3984454	71.3663158	0.7482304	NM_014062	NOB1P	
0.0021903	233.0709896	183.0490111	1.273271	NM_001031677	RAB24	
0.002331	78.1171224	97.3888835	0.8021154	NM_017699	SIDT1	
0.002457	55.4244192	75.0410081	0.7385884	NM_020642	C11orf17	
0.0025345	956.4176519	673.3678999	1.4203493	NM_033405	PRIC285	
0.002748	111.8853999	65.758725	1.7014533	NM_015071	ARHGAP26	
0.002797	77.7048739	43.8710092	1.7712124	NM_018984	SSH1	Regulation of actin cytoskeleton
0.0031593	96.3060504	70.9280275	1.3577996	NM_001031719	FLJ13639	
0.0032597	74.3860187	107.9506798	0.6890741	NM_006134	TMEM50B	
0.0034606	144.2016183	198.6110208	0.7260504	NM_144638	TMEM42	Role of Ran in mitotic spindle regulation
0.0035429	121.5756396	65.759618	1.8487887	NM_052828	TRIM10	
0.0035783	76.7861621	100.233922	0.7660696	NM_181340	WDR21A	
0.0037429	1428.384827	1074.9697	1.3287675	NM_006573	TNFSF13B	TACI and BCMA stimulation of B cell immune responses., Cytokine-cytokine receptor interaction
0.0038303	2308.950828	1401.307414	1.6477118	NM_003494	DYSF	
0.003983	287.009692	225.0477753	1.2753278	NM_001776	ENTPD1	
0.004009	136.1716197	185.1636214	0.7354124	NM_006985	NPIP	
0.0040379	89.442346	60.228624	1.4850471	NM_002463	MX2	cell_signaling
0.0041087	636.3160782	820.7957534	0.7752429	NM_024619	FN3KRP	
0.0041129	54.5849904	70.7770909	0.771224	NM_024493	ZNF306	
0.0042	129.564397	206.5903427	0.6271561	NM_024048	MGC3020	
0.0043066	77.2854192	113.9056458	0.6785039	NM_004284	CHD1L	
0.0044582	313.8851602	213.5519415	1.4698305	NM_000167	GK	Glycerolipid metabolism, PPAR signaling pathway
0.004488	106.5122966	151.1224528	0.7048079	NM_018346	RSAD1	
0.0045615	64.1665543	96.0495151	0.668057	NM_138341	TMEM116	
0.0051912	433.4919731	308.0187609	1.4073557	NM_002093	GSK3B	
0.0053976	83.1198159	48.5045894	1.7136485	NM_203411	TMEM88	
0.0054644	249.0470921	304.1155458	0.8189226	NM_020444	KIAA1191	
0.0057117	1224.60185	822.4718523	1.4889286	NM_022059	CXCL16	
0.0057841	110.4525818	142.0882007	0.7773522	NM_080605	B3GALT6	Chondroitin sulfate biosynthesis, Glycan structures - biosynthesis 1
0.0059993	6755.193466	4676.588102	1.4444705	NM_006931	SLC2A3	Vitamin C in the Brain, cell_signaling, misc, pharmacology
0.0060617	57.8154193	37.7067951	1.5332891	NM_020439	CAMK1G	
0.0062033	107.4822883	141.3133872	0.7605952	NM_024691	ZNF419	
0.0062233	393.164908	285.6590853	1.3763431	NM_003268	TLR5	Toll-like receptor signaling pathway, immunology
0.006241	706.5141744	366.4292943	1.9281051	NM_000717	CA4	Nitrogen metabolism
0.0062475	4308.608064	2917.50243	1.4768139	NM_021642	FCGR2A	immunology
0.0063598	1500.259142	1159.001672	1.2944409	NM_004031	IRF7	gene_regulation, transcription
0.0064295	4468.985126	3070.56217	1.455429	NM_012387	PADI4	
0.0065349	162.9657928	206.0056073	0.7910745	NM_012139	SERGEF	
0.0065475	3763.198465	2463.563116	1.5275429	NM_022083	C1orf24	
0.0065818	169.8341703	202.8024016	0.8374367	NM_002857	PEX19	
0.0068421	121.5985049	67.4619564	1.8024752	NM_130385	MRVI1	

Personal non-commercial use only. The Journal of Rheumatology Copyright © 2010. All rights reserved.

0.0069067	4094.439362	3127.288913	1.3092616	NM_139266	STAT1	Apoptotic Signaling in Response to DNA Damage, EGF Signaling Pathway, IFN alpha signaling pathway, IFN gamma signaling pathway, IL22 Soluble Receptor Signaling Pathway, Inhibition of Cellular Proliferation by Gleevec, MAPKinase Signaling Pathway, p38 MAPK Signaling Pathway, PDGF Signaling Pathway, TPO Signaling Pathway, Jak-STAT signaling pathway, Toll-like receptor signaling pathway, angiogenesis, immunology, signal_transduction
0.0069214	79.185264	50.4842138	1.5685153	NM_004347	CASP5	MAPK signaling pathway, apoptosis, immunology
0.0069767	2792.841513	2234.522646	1.2498605	NM_033306	CASP4	Caspase Cascade in Apoptosis, MAPK signaling pathway, apoptosis, immunology
0.0069791	57.8324274	41.4198501	1.3962491	NM_024496	C14orf4	
0.0070874	458.6254977	334.7400414	1.3700945	NM_032536	NTNG2	
0.0072365	309.1783279	239.2651331	1.2921997	NM_080424	SP110	
0.0075236	2360.736625	1641.346746	1.4382924	NM_001995	ACSL1	Perou's- Intrinsic- Breast-Cancer-Genes, Adipocytokine signaling pathway, Fatty acid metabolism, PPAR signaling pathway
0.0075429	101.3066589	143.8378404	0.7043116	NM_016311	ATPIF1	
0.0075842	55.9221565	71.0356931	0.7872402	NM_002913	RFC1	
0.007629	284.5814239	197.0116273	1.4444905	NM_138557	TLR4	immunology
0.0078644	8192.608896	5797.077578	1.4132309	NM_020980	AQP9	
0.0079295	352.6683885	228.2252792	1.5452644	NM_002243	KCNJ15	
0.0079915	69.4960341	55.0535557	1.2623351	NM_171846	LACTB	
0.0080308	4230.418121	2883.942998	1.4668869	NM_022083	C1orf24	
0.0080335	61.6309441	80.2352561	0.768128	NM_001032999	CBFA2T2	cell_cycle, immunology
0.0080701	78.1569398	53.726485	1.454719	NM_001924	GADD45A	ATM Signaling Pathway, Cell Cycle: G2/M Checkpoint, Hypoxia and p53 in the Cardiovascular system, p53 Signaling Pathway, Cell cycle, MAPK signaling pathway
0.0083262	1358.69924	985.1936551	1.379119	NM_030769	NPL	
0.0084262	961.6377224	682.6056177	1.408775	NM_006825	CKAP4	
0.0084503	87.9484113	65.2853476	1.3471386	NM_138962	MSI2	
0.0085554	597.6472872	398.0354876	1.5014925	NM_005306	FFAR2	
0.0087672	95.5304166	51.085383	1.8700147	NM_000062	SERPING1	Intrinsic Prothrombin Activation Pathway, Complement and coagulation cascades
0.0090354	158.4764733	192.559439	0.8230003	NM_004793	PRSS15	
0.0091097	198.4911419	136.4177609	1.4550242	NM_144653	BTBD14A	
0.009238	83.129073	110.3058286	0.7536236	NM_017607	PPP1R12C	
0.0092556	126.8566834	181.5263622	0.6988334	NM_139281	WDR36	
0.0092918	70.2149582	85.965983	0.8167761	NM_014149	HSPC049	
0.0093926	117.3907998	154.3682162	0.7604597	NM_017541	CRYGS	
0.0095282	154.6194436	194.177994	0.7962769	NM_024056	TMEM106C	
0.0095422	122.7966001	150.2197121	0.8174466	NM_001002880	PGEA1	
0.0095887	892.9376363	1196.945428	0.7460137	NM_019063	EML4	
0.0096633	89.6226922	56.5656362	1.5844017	NM_017409	HOXC10	development
0.0096843	85.2460615	116.6847675	0.7305672	NM_024070	MGC2463	
0.0097158	114.2832903	153.2233268	0.7458609	NM_206930	SYTL2	
0.0097279	59.7622428	38.6757469	1.5452124	NM_000208	INSR	Control of skeletal myogenesis by HDAC & calcium/calmodulin-dependent kinase (CaMK), Growth Hormone Signaling Pathway, Insulin Signaling Pathway, Adherens junction, Dentatorubropallidoluyian atrophy (DRPLA), Insulin signaling pathway, Type II diabetes mellitus, immunology

oligo- and polyarticular, resembles spondyloarthritis more than it does rheumatoid arthritis. *Arthritis Res Ther* 2005;7:R569-80.

34. Baeten D, Demetter P, Cuvelier CA, Kruithof E, Van Damme N, De Vos M, et al. Macrophages expressing the scavenger receptor

CD163: a link between immune alterations of the gut and synovial inflammation in spondyloarthritis. *J Pathol* 2002;196:343-50.

35. Weaver LK, Pioli PA, Wardwell K, Vogel SN, Guyre PM. Up-regulation of human monocyte CD163 upon activation of

Appendix 2. Genes that were only differentially expressed in AS.

Parametric p value	Geom mean of intensities in class 1 (AS)	Geom mean of intensities in class 2 (control)	Fold-change	GB acc	Gene symbol	Defined Gene list	Gene involved in the dysregulated pathways in Table 3
0.0006957	9364.31931	4899.47231	1.9112914	NM_000570	FCGR3B	Natural killer cell mediated cytotoxicity, immunology	Dendritic Cell Maturation
0.0009312	70.135071	39.2769027	1.7856569	NM_198330	INPP5F		
0.0011141	52.8656567	70.9770793	0.7448272	NM_001013840	C10orf61		
0.0017355	61.4036443	44.5498863	1.378312	NM_001125	ADPRH		
0.0018922	87.5243114	47.5085126	1.8422869	NM_170606	MLL3	Lysine degradation	
0.002143	53.3984454	71.3663158	0.7482304	NM_014062	NOB1P		
0.0021903	233.07099	183.049011	1.273271	NM_001031677	RAB24		
0.002331	78.1171224	97.3888835	0.8021154	NM_017699	SIDT1		
0.002457	55.4244192	75.0410081	0.7385884	NM_020642	C11orf17		
0.002748	111.8854	65.758725	1.7014533	NM_015071	ARHGAP26		
0.002797	77.7048739	43.8710092	1.7712124	NM_018984	SSH1	Regulation of actin cytoskeleton	
0.0031593	96.3060504	70.9280275	1.3577996	NM_001031719	FLJ13639		
0.0035429	121.57564	65.759618	1.8487887	NM_052828	TRIM10		
0.0035783	76.7861621	100.233922	0.7660696	NM_181340	WDR21A		
0.0038303	2308.95083	1401.30741	1.6477118	NM_003494	DYSF		
0.0041087	636.316078	820.795753	0.7752429	NM_024619	FN3KRP		
0.0041129	54.5849904	70.7770909	0.771224	NM_024493	ZNF306		
0.0043066	77.2854192	113.905646	0.6785039	NM_004284	CHD1L		
0.0044582	313.88516	213.551942	1.4698305	NM_000167	GK	Glycerolipid metabolism, PPAR signaling pathway	
0.004488	106.512297	151.122453	0.7048079	NM_018346	RSAD1		
0.0045615	64.1665543	96.0495151	0.668057	NM_138341	TMEM116		
0.0051912	433.491973	308.018761	1.4073557	NM_002093	GSK3B		
0.0053976	83.1198159	48.5045894	1.7136485	NM_203411	TMEM88		
0.0057841	110.452582	142.088201	0.7773522	NM_080605	B3GALT6	Chondroitin sulfate biosynthesis, Glycan structures - biosynthesis 1	
0.0059993	6755.19347	4676.5881	1.4444705	NM_006931	SLC2A3	Vitamin C in the Brain, cell_signaling, misc, pharmacology	
0.0062233	393.164908	285.659085	1.3763431	NM_003268	TLR5	Toll-like receptor signaling pathway, immunology	NF-κB Signaling, Toll-like Receptor Signaling, TREM1 Signaling
0.006241	706.514174	366.429294	1.9281051	NM_000717	CA4	Nitrogen metabolism	
0.0062475	4308.60806	2917.50243	1.4768139	NM_021642	FCGR2A	immunology	Dendritic Cell Maturation
0.0064295	4468.98513	3070.56217	1.455429	NM_012387	PADI4		
0.0065349	162.965793	206.005607	0.7910745	NM_012139	SERGEF		

cell-surface Toll-like receptors. *J Leukoc Biol* 2007;81:663-71.

36. Jiang W, Sun R, Wei H, Tian Z. Toll-like receptor 3 ligand attenuates LPS-induced liver injury by down-regulation of toll-like receptor 4 expression on macrophages. *Proc Natl Acad Sci USA* 2005;102:17077-82.

37. Fert I, Glatigny S, Poulain C, Satumtira N, Dorris ML, Taurog JD, et al. Correlation between dendritic cell functional defect and spondylarthritis phenotypes in HLA-B27/Human beta2-microglobulin-transgenic rat lines. *Arthritis Rheum* 2008;58:3425-9.

0.0065349	162.965793	206.005607	0.7910745	NM_012139	SERGEF		
0.0065475	3763.19846	2463.56312	1.5275429	NM_022083	C1orf24		
0.0065818	169.83417	202.802402	0.8374367	NM_002857	PEX19		
0.0068421	121.598505	67.4619564	1.8024752	NM_130385	MRV11		
0.0069791	57.8324274	41.4198501	1.3962491	NM_024496	C14orf4		
0.0075236	2360.73663	1641.34675	1.4382924	NM_001995	ACSL1	Perou's- Intrinsic- Breast-Cancer-Genes, Adipocytokine signaling pathway, Fatty acid metabolism, PPAR signaling pathway	
0.0075429	101.306659	143.83784	0.7043116	NM_016311	ATPIF1		
0.0075842	55.9221565	71.0356931	0.7872402	NM_002913	RFC1		BRCA1 in DNA Damage Response
0.007629	284.581424	197.011627	1.4444905	NM_138557	TLR4	immunology	NF-κB Signaling, Dendritic Cell Maturation, Toll-like Receptor Signaling, TREM1 Signaling
0.0078644	8192.6089	5797.07758	1.4132309	NM_020980	AQP9		
0.0079295	352.668389	228.225279	1.5452644	NM_002243	KCNJ15		
0.0079915	69.4960341	55.0535557	1.2623351	NM_171846	LACTB		
0.0080308	4230.41812	2883.943	1.4668869	NM_022083	C1orf24		
0.0080701	78.1569398	53.726485	1.454719	NM_001924	GADD45A	ATM Signaling Pathway, Cell Cycle: G2/M Checkpoint, Hypoxia and p53 in the Cardiovascular system, p53 Signaling Pathway, Cell cycle, MAPK signaling pathway	BRCA1 in DNA Damage Response
0.0083262	1358.69924	985.193655	1.379119	NM_030769	NPL		
0.0090354	158.476473	192.559439	0.8230003	NM_004793	PRSS15		
0.0091097	198.491142	136.417761	1.4550242	NM_144653	BTBD14A		
0.009238	83.129073	110.305829	0.7536236	NM_017607	PPP1R12C		
0.0092556	126.856683	181.526362	0.6988334	NM_139281	WDR36		
0.0095282	154.619444	194.177994	0.7962769	NM_024056	TMEM106C		
0.0095887	892.937636	1196.94543	0.7460137	NM_019063	EML4		
0.0096633	89.6226922	56.5656362	1.5844017	NM_017409	HOXC10	development	
0.0097279	59.7622428	38.6757469	1.5452124	NM_000208	INSR	Control of skeletal myogenesis by HDAC & calcium/calmodulin-dependent kinase (CaMK), Growth Hormone Signaling Pathway, Insulin Signaling Pathway, Adherens junction, Dentatorubropallidolusian atrophy (DRPLA), Insulin signaling pathway, Type II diabetes mellitus, immunology	NF-κB Signaling